

Common Deficiencies-OGD Considerations

Complex Generic Drug Product Development Workshop

How to resolve current challenges in ANDAs for topical products

September 13, 2018

Usha Katragadda

Division of Bioequivalence III, Office of Bioequivalence

Office of Generic Drugs

CDER | US FDA

Disclaimer



- The views expressed in this presentation do not reflect the official policies of the FDA, or the Department of Health and Human Services; nor does any mention of trade names, commercial practices, or organization imply endorsement by the United States Government.
- I do not have any financial interest or conflict of interest with any pharmaceutical companies.

Product Specific Guidances: Semisolid Topical Products



- Acyclovir Ointment, Silver Sulfadiazine Cream and Docosanol Cream : Q1/Q2/Q3 + **IVRT**
- Acyclovir Cream: Q1/Q2/Q3 + **IVRT** + **IVPT**
- Benzyl Alcohol Lotion: Q1/Q2/Q3 + **IVRT** + Lice Assay
- Dapsone Gel and Ivermectin Cream : Q1/Q2/Q3 + **IVRT** + **IVPT** +
In-Vivo pK BE Study

In Vitro Release Test (IVRT)



Conducting an IVRT:

As described in the United States Pharmacopeia (USP) General Chapter <1724>, Semisolid Drug Products – Performance Tests.

IVRT

What should be submitted for evaluation?

- IVRT Method Development Report
- IVRT Method Validation Report
- Analytical Method validation Report
- IVRT Pivotal Study Report
- Standard Operating Procedure (SOP)

IVRT Method Development



Method Parameters:

Product dose amount, Stirring rate, Sampling times, IVRT apparatus


Membrane: Inertness

Receptor Solution: Solubility and stability

IVRT Method Development



Membrane Evaluation




IVRT Method Validation



Validation Components



- Linearity and Range
- Accuracy/Precision and Reproducibility
- Recovery, Mass Balance & Dose Depletion



IVRT Method Validation

Sensitivity and selectivity

Specificity (Proportionality)



IVRT Method Validation

Validation Components

- Robustness
- Membrane Inertness
- Receptor Solution Solubility
- Apparatus Qualification

Deficiencies for IVRT studies



Experimental Details:

- Stir speed
- Membrane pre-soak time
- Degassing of the receptor medium
- Sample withdrawal
- Amount of sample loaded (dose per cell)

Deficiencies for IVRT studies



Data Analysis:

- Report on data analysis
- Raw numerical data : Dilution factor, calibration standards, and quality control samples
- IVRT data in SAS Transport format

Deficiencies for IVRT studies

Incomplete Study Reports : Analytical and IVRT

- Method development reports
- Method validation reports
- Standard Operating Procedure (SOP)

In Vitro Permeation Test (IVPT)



Conducting an IVPT:

Using excised human skin mounted on a qualified diffusion cell system.

IVPT




What should be submitted for evaluation?

- IVPT Method Development Report
- IVPT Method Validation Report
- Analytical Method validation Report
- IVPT Pilot Study Report
- IVPT Pivotal Study Report
- Standard Operating Procedure (SOP)

IVPT Method Development



Method Parameters: *Product dose amount, Stirring rate, Sampling times*



IVPT Method Validation

Validation Components

- Membrane (skin) qualification
- Receptor Solution Qualification
- Receptor Solution Sampling Qualification
- Receptor Solution Sample Analytical Method Validation
- Environmental Control

IVPT Method Validation

Validation Components

- Permeation Profile and Range
- Precision and Reproducibility
- Recovery, Mass Balance & Dose Depletion
- Discrimination Sensitivity and Selectivity
- Robustness

IVPT Pilot Study

- To estimate number of donors required for the pivotal study.
- Multiple skin donors: minimum of 4 replicate skin sections per donor per treatment group.
- Parallel assessment should be performed with a third product.
- Results: should not be combined with the IVPT pivotal study.

IVPT Pivotal Study

- Non-dosed control skin section from each donor
- Pre-dose zero sample from each diffusion cells
- Duration of an IVPT study: maximum flux and a decline in the flux
- Dose staggering and sampling synchronization

Deficiencies for IVPT studies



Study Design:

- No control skin section (non dosed)
- Insufficient sampling times
- Skin from only single donor
- Skin donor Inclusion/Exclusion criteria not submitted
- Randomization sequence not submitted
- No rationale for clinically irrelevant dose

Deficiencies for IVPT studies



Experimental Details:

- Temperature
- Size of the tissue mounted on the Franz diffusion cell
- Skin conditions (occluded vs unoccluded)
- Speed of rotation (stir bar)
- Dosing and sampling design

Deficiencies for IVPT studies



Data Analysis:

- Skin sections
- Raw numerical data : Dilution factor, calibration standards, and quality control samples
- IVPT data in SAS Transport format

Deficiencies for IVPT studies

Incomplete Study Reports : Analytical and IVPT

- Method development reports
- Method validation reports
- Standard Operating Procedure (SOP)

Summary



- IVRT/IVPT data reviewed by Division of Bioequivalence.
- The method validation should be performed using validated sample analytical procedures under principles of GLP.
- Inadequate submission of the data may often lead to the delay in the review process and the final approval of the application.

Acknowledgements



Office of Bioequivalence

- Dale Conner, PharmD
- Trueman Sharp, M.D.
- Nilufer Tampal, Ph.D
- April Braddy, Ph.D
- Ke Ren, Ph.D

OGD Office of Research and Standards

- Sam Raney, Ph.D.
- Priyanka Gosh

